I 49.89:82/47-1/map C18

MMS 85-0011
FWS/OBS-82/47.1

MAP C18

# Tarpon Springs

## FLORIDA ECOLOGICAL ATLAS

### SOILS AND LANDFORMS

TARPON SPRINGS, FLORIDA

30 X 60 MINUTE SERIES (TOPOGRAPHIC)




ORIGINAL

Scale 1:100 000

TOPICAL MAPS IN THIS ATLAS:
A. BIOLOGICAL RESOURCES
B. SOCIOECONOMIC FEATURES
C. SOILS AND LANDFORMS
D. OIL, GAS AND MINERALS
E. HYDROLOGY AND CLIMATOLOGY

1984

Produced by U.S. Fish and Wildlife Service and Minerals Management Service

Projection and 10,000 meter grid, zone 17: Universal Transverse Mercator


GULF OF MEXICO
Anclote Anchorage
Anclote Keys
PASCO CO
HILLSBOROUGH CO
Tarpon Springs
Port Richey
New Port Richey
Holiday
Hudson
Elfers
Odessa
Lutz
Land O' Lakes
Zephyrhills
Dade City
Crystal Springs
Wesley Chapel
Trilby
Starkey
Seven Oaks
Saddle Creek
Lacoochee
Trilby
Pithlachascotee
Anclote
Cotee
Big Cypress
Hillsborough
Alligator Lake
Crystal Lake
Lake Padgett
Lake Tarpon
Lake Pasadena
Lake Seminole
Lake Thonotosassa
Lake Magdalene
Lake Carroll
Lake Saunders
Lake Bernadette
Lake Starkey
Lake Patience
Lake Jovita
Lake Iola
Lake Fern
Lake Myrtle
Lake Lucy
Lake Linda
Lake Lizzie
Lake Lindsey
Lake Lashbrook
Lake Jessamine
Lake Hanna
Lake Hancock
Lake Gall
Lake Fannie
Lake Eloise
Lake Deeson
Lake Crenshaw
Lake Bonnet
Lake Bonny
Lake Bird
Lake Bethel
Lake Armistead
Lake Alice
Lake Agnes
Lake Ada
Lake Abbott
Lake Abbey
Lake Abbie
Lake Abbe
Lake Abbe
Lake Abbe


PINELLAS COUNTY
AQUATIC PRESERVE
PASCO CO
HILLSBOROUGH CO
TAMPA
Caladesi Island
Honeymoon Island
Crystal Beach
Palm Harbor
Dunedin
Clearwater
Safety Harbor
Oldsmar
Tarpon Springs
Hillsborough River
Lake Thonotosassa
Lake Roberts
Lake Park
Lake Magdalene
Lake Tampa
Lake Chapman
Lake Cooper
Lake Fern
Lake Forest
Lake Ellen
Lake Lutz
Lake Lizzie
Lake Ida
Lake Bird
Lake Hanna
Lake Keene
Lake Mable
Lake Mary
Lake Ann
Lake Jackson
Lake Hillsborough
Lake Thomas
Lake Crenshaw
Lake Seminole
Lake Tarpon
Lake Butler
Lake Brant
Lake Haines
Lake Lindsey
Lake Ponce
Lake Fant
Lake Lucy
Lake Ashley
Lake Hunter
Lake Hooker
Lake Hobbs
Lake Hilda
Lake Hilma
Lake Hilton
Lake Hiawassee
Lake Hiawatha
Lake Hiawatha
Lake Hiawatha

SCALE 1:100 000

1 CENTIMETER ON THE MAP REPRESENTS 1 KILOMETER ON THE GROUND.
CONTOUR INTERVAL 5 METERS

KILOMETERS 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

MILES 0 1 2 3 4 5 6 7 8 9 10 11 12

0 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000 11000 12000 13000 14000 15000 16000 17000 18000 19000 20000 METERS

0 5000 10000 15000 20000 25000 30000 35000 40000 45000 50000 55000 60000 65000 70000 FEET

# TARPON SPRINGS, FLA.

N2800—W8200/30 X 60

1978

THE FLORIDA ECOLOGICAL ATLAS WAS PRODUCED BY THE U.S. FISH AND WILDLIFE SERVICE, DIVISION OF BIOLOGICAL SERVICES, NATIONAL COASTAL ECOSYSTEMS TEAM (FWS 1280). BC 470 AND THE MINERALS MANAGEMENT SERVICE FOR COASTAL ECOSYSTEMS INFORMATION NEEDED TO ASSESS THE POTENTIAL IMPACTS AND EVALUATE CHANGE AND TO RESOLVE CONFLICTS ARISING LAND USES, FISH AND WILDLIFE RESOURCES AND OUTER CONTINENTAL SHELF OIL AND GAS DEVELOPMENT. THE STUDY AREA INCLUDES THE COASTAL COUNTIES ESCAMBIA, SANTA ROSA, OKALOOSA, WALTON, BAY, GULF, FRANKLIN, WAKULLA, JEFFERSON, TAYLOR, DIXIE, LEVY, CITRUS AND HERNANDO, AND EXTENDS OFFSHORE APPROXIMATELY 10 MILES. THERE IS NO ATTEMPT IN THE DESIGN OF THIS ATLAS TO DELINEATE OR DEFINE THE EXACT OFFSHORE LIMITS OF JURISDICTION OF ANY FEDERAL, STATE, OR LOCAL GOVERNMENT TO ESTABLISH THE GEOGRAPHICAL SCOPE OF THE REGULATORY PROGRAMS OF GOVERNMENT AGENCIES.

THIS MAP HAS BEEN COMPILED FROM A VARIETY OF SOURCE MATERIALS WITH VARYING DEGREES OF ACCURACY. DEVIATIONS MAY BE PRESENT WHICH CANNOT BE DETECTED. USERS OF RESOURCES AND OTHER MAPPED DATA ARE CAUTIONED TO CONSULT PRIMARY SOURCES WHERE MAPS FOR PRECISE MEASUREMENTS OF DISTANCES OR AREAS.

## PRIMARY SOURCES OF MAPPED INFORMATION:

LANDFORMS: Physiographic Map of Florida, State Bureau of Geology, 1960.

SOILS: General Soils Atlas of Florida, Division of State Planning, 1975. Soil Survey Reports, 1958-1978, Soil Conservation Service, 1982.

ACTIVE DUNES: Black and White Aerial Photographs, 1:80,000 Scale, Mark Hurd Aerial Surveys, Inc., 1982. Topographic Maps, 1:24,000 Scale, U.S. Geological Survey, various dates.

GEOLOGIC FAULTS: Map of Escambia and Santa Rosa Counties, Florida, and Adjacent Parts of the Coastal Complex of the Florida Panhandle, Hayden, State Bureau of Geology, 1980.

FOR COMPLETE REFERENCES SEE MAP NARRATIVE

66. Candler-Pedro-Tavares Association: Nearly level to strongly sloping, poorly drained fine sands and sandy loams overlying loamy and clayey material.

68. Arredondo-Kendrick Association: Nearly level to sloping, well drained soils with very thick sandy layers over thin loamy or sandy loam lamellae and occasionally and moderately well drained soils, sands throughout.

71. Myakka-Basinger Association: Nearly level, poorly drained and very poorly drained soils, sands subsoil and ponds drained soils, sands throughout.

74. Arredondo-Sparr-Kendrick Association: Nearly level, to sloping, well drained and somewhat poorly drained sandy soils over loamy material.

76. Wauseon Association: Nearly level to gently sloping, moderately well drained soils overlying loamy material.

77. Nobleton-Bichton-Flemington Association: Nearly level to strongly sloping, poorly drained fine sands and sandy loams overlying loamy and clayey material.

78. EauGallie-Wabasso-Basinger Association: Nearly level, poorly drained sandy soils which are either sands throughout or overlie loamy material.

79. Pomello-Floridana-Wabasso Association: Nearly level, poorly drained sandy soils overlying a clayey subsoil or loamy material.

80. Okeelanta-Arapaha-Terra Ceia Association: Nearly level, very poorly drained organic and mineral soils overlying sandy material or loamy soils overlying limestone.

81. Homosassa-Weekiwachee-Lacoochee Association: Nearly level, very poorly drained organic and mineral soils, and poorly drained, thin, sandy soils overlying limestone which are subject to frequent tidal flooding.

83. Salt Water Marsh and Swamp Association: Nearly level, very poorly drained soils subject to frequent tidal flooding.

84. Fresh Water Swamp and Marsh Association: Nearly level, very poorly drained soils subject to prolonged flooding.

85. Bichton-Lochloosa-Kendrick Association: Nearly level to sloping, poorly drained sandy soils with loamy subsoils, somewhat poorly drained soils with thick sandy layers over loamy subsoil, and well drained sands with very thick sandy layers over loamy subsoil.

88. Pomello-Sunderland Association: Nearly level to gently sloping, somewhat poorly drained and well drained soils with thin sandy layers over clayey subsoil.

91. Myakka-Pomello-Basinger Association: Nearly level, poorly drained and moderately well drained sandy soils with weakly cemented sandy subsoil and poorly drained soils sandy throughout.

E = ft/yr — BEACH EROSION AREA (Period of Record: 1957-1981; Univ. of West Florida, 1981)

A = ft/yr — BEACH ACCRETION AREA (Period of Record: 1957-1981; Univ. of West Florida, 1981)

GULF BARRIER ISLAND

ACTIVE DUNE AREA

U D — NORMAL FAULT (Showing relative vertical movement, U. up, and D. down)

U D — SUB-SURFACE NORMAL FAULT (Showing relative vertical movement, U. up, and D. down)

STUDY AREA BOUNDARY (Offshore)

Inland (County)

Inland (State)


84°00'
81°00'
30°00'
27°30'
ESCAMBIA
SANTA ROSA
OKALOOSA
WALTON
BAY
GULF
FRANKLIN
WAKULLA
JEFFERSON
TAYLOR
DIXIE
LEVY
CITRUS
HERNANDO
PASCO
PINELLAS
HILLSBOROUGH
MANATEE
SARASOTA
CHARLOTTE
LEE
COLLIER
MONROE
DADE
BROWARD
PALM BEACH
MARTIN
ST. LUCIE
INDIAN RIVER
OKEECHOBEE
HENDRY
GLADES
BROWARD
DADE
MONROE
COLLIER
LEE
CHARLOTTE
MANATEE
SARASOTA
DESOTO
HARDEE
HIGHLANDS
OKEECHOBEE
INDIAN RIVER
ST. LUCIE
MARTIN
PALM BEACH
OKEECHOBEE
GLADES
HENDRY
OSCEOLA
POLK
PINELLAS
PASCO
HERNANDO
SUMTER
LAKE
ORANGE
SEMINOLE
VOLUSIA
BREVARD
OSCEOLA
POLK
HARDEE
DESOTO